Contents:

Volume 134, Number 1, July 1996

1 ***In situ* analysis of changes in telomere size during replicative aging and cell transformation.**
S. Henderson, R. Allsopp, D. Spector, S.-S. Wang and C. Harley

13 **Bcl-2 down-regulates the activity of transcription factor NF-κB induced upon apoptosis.**
S. Grimm, M.K.A. Bauer, P.A. Baeuerle, and K. Schulze-Osthoff

25 **Signal sequence–dependent function of the TRAM protein during early phases of protein transport across the endoplasmic reticulum membrane.**
S. Voigt, B. Jungnickel, E. Hartmann, and T.A. Rapoport

37 **The absence of Pmp47, a putative yeast peroxisomal transporter, causes a defect in transport and folding of a specific matrix enzyme.**
Y. Sakai, A. Saiganji, H. Yurimoto, K. Takabe, H. Saiki, and N. Kato

53 **Myosin filament structure in vertebrate smooth muscle.**
J.-Q. Xu, B.A. Harder, P. Uman, and R. Craig

67 **A function for filamentous α-smooth muscle actin: Retardation of motility in fibroblasts.**
L. Rønnov-Jessen and O.W. Petersen

81 ***Drosophila* paramyosin/miniparamyosin gene products show a large diversity in quantity, localization, and isoform pattern: A possible role in muscle maturation and function.**
M. Maroto, J. Arredondo, D. Goulding, R. Marco, B. Bullard, and M. Cervera

93 **Forced expression of dystrophin deletion constructs reveals structure–function correlations.**
J.A. Rafael, G.A. Cox, K. Corrado, D. Jung, K.P. Campbell, and J.S. Chamberlain

103 **Live analysis of free centrosomes in normal and aphidicolin-treated *Drosophila* embryos.**
A. Debec, R.F. Kalpin, D.R. Daily, P.D. McCallum, W.F. Rothwell, and W. Sullivan

117 **The *Saccharomyces cerevisiae* actin-related protein Arp2 is involved in the actin cytoskeleton.**
V. Moreau, A. Madania, R.P. Martin, and B. Winsor

133 **Armadillo is required for adherens junction assembly, cell polarity, and morphogenesis during *Drosophila* embryogenesis.**
R.T. Cox, C. Kirkpatrick, and M. Peifer

149 ***armadillo*, *bazooka*, and *stardust* are critical for early stages in formation of the zonula adherens and maintenance of the polarized blastoderm epithelium in *Drosophila*.**
H.-A.J. Müller and E. Wieschaus

165 **The adenomatous polyposis coli tumor suppressor protein localizes to plasma membrane sites involved in active cell migration.**
I.S. Näthke, C.L. Adams, P. Polakis, J.H. Sellin, and W.J. Nelson.

181 **SPC4, SPC6, and the novel protease SPC7 are coexpressed with bone morphogenetic proteins at distinct sites during embryogenesis.**
D.B. Constam, M. Calfon, and E.J. Robertson

193 **Interference with p53 protein inhibits hematopoietic and muscle differentiation.**
S. Soddu, G. Blandino, R. Scardigli, S. Coen, A. Marchetti, M.G. Rizzo, G. Bossi, L. Cimino, M. Crescenzi, and A. Sacchi

205 **Growth cone collapse and inhibition of neurite growth by *botulinum* neurotoxin C1: A t-SNARE is involved in axonal growth.**
M. Igarashi, S. Kozaki, S. Terakawa, S. Kawano, C. Ide, and Y. Komiya

217 **Intracellular signals direct integrin localization to sites of function in embryonic muscles.**
M.D. Martin-Bermudo and N.H. Brown

227 ***Xenopus* embryonic cell adhesion to fibronectin: Position-specific activation of RGD/synergy site–dependent migratory behavior at gastrulation.**
J.W. Ramos and D.W. DeSimone

241 **The intracellular functions of $\alpha_6\beta_4$ integrin are regulated by EGF.**
F. Mainiero, A. Pepe, M. Yeon, Y. Ren, and F.G. Giancotti

255 **Biology of chemokine and classical chemoattractant receptors: Differential requirements for adhesion-triggering versus chemotactic responses in lymphoid cells.**
J.J. Campbell, S. Qin, K.B. Bacon, C.R. Mackay, and E.C. Butcher

267 ADDITIONS AND CORRECTIONS

Volume 134, Number 2, July 1996

269 **Signal sequences specify the targeting route to the endoplasmic reticulum membrane.**
D.T.W. Ng, J.D. Brown, and P. Walter

279 **Endoplasmic reticulum localization of Sec12p is achieved by two mechanisms: Rer1p-dependent retrieval that requires the transmembrane domain and Rer1p-independent retention that involves the cytoplasmic domain.**
M. Sato, K. Sato, and A. Nakano

295 **Evidence that phospholipase D mediates ADP ribosylation factor–dependent formation of Golgi coated vesicles.**
N.T. Ktistakis, H.A. Brown, M.G. Waters, P.C. Sternweis, and M.G. Roth

307 **The human mitochondrial import receptor, hTom20p, prevents a cryptic matrix targeting sequence from gaining access to the protein translocation machinery.**
H.M. McBride, I.S. Goping, and G.C. Shore

315 **Two components of the chloroplast protein import apparatus, IAP86 and IAP75, interact with the transit sequence during the recognition and translocation of precursor proteins at the outer envelope.**
Y. Ma, A. Kouranov, S.E. LaSala, and D.J. Schnell

329 **Poisson-distributed active fusion complexes underlie the control of the rate and extent of exocytosis by calcium.**
S.S. Vogel, P.S. Blank, and J. Zimmerberg

339 **Endocytosis of chimeric influenza virus hemagglutinin proteins that lack a cytoplasmic recognition feature for coated pits.**
J. Lazarovits, H.Y. Naim, A.C. Rodriguez, R.-H. Wang, E. Fire, C. Bird, Y.I. Henis, and M.G. Roth

349 **Host cell invasion by trypanosomes requires lysosomes and microtubule/kinesin–mediated transport.**
A. Rodríguez, E. Samoff, M.G. Rioult, A. Chung, and N.W. Andrews

363 **Sphingolipids and glycoproteins are differentially trafficked to the *Chlamydia trachomatis* inclusion.**
M.A. Scidmore, E.R. Fischer, and T. Hackstadt

375 **Formation of triads without the dihydropyridine receptor α subunits in cell lines from dysgenic skeletal muscle.**
J.A. Powell, L. Petherbridge, and B.E. Flucher

389 **Coordinated regulation of platelet actin filament barbed ends by gelsolin and capping protein.**
K. Barkalow, W. Witke, D.J. Kwiatkowski, and J.H. Hartwig

401 **Smoothelin, a novel cytoskeletal protein specific for smooth muscle cells.**
F.T.L. van der Loop, G. Schaart, E.D.J. Timmer, F.C.S. Ramaekers, and G.J.J.M. van Eys

413 **The bud4 protein of yeast, required for axial budding, is localized to the mother/bud neck in a cell cycle–dependent manner.**
S.L. Sanders and I. Herskowitz

429 **γ-Tubulin–like Tub4p of *Saccharomyces cerevisiae* is associated with the spindle pole body substructures that organize microtubules and is required for mitotic spindle formation.**
A. Spang, S. Geissler, K. Grein, and E. Schiebel

443 **Analysis of Tub4p, a yeast γ-tubulin-like protein: Implications for microtubule-organizing center function.**
L.G. Marschall, R.L. Jeng, J. Mulholland, and T. Stearns

455 **Anastral meiotic spindle morphogenesis: Role of the Non-Claret Disjunctional kinesin-like protein.**
H.J.G. Matthies, H.B. McDonald, L.S.B. Goldstein, and W.E. Theurkauf

465 **ARIA/HRG regulates AChR ε subunit gene expression at the neuromuscular synapse via activation of phosphatidylinositol 3-kinase and Ras/MAPK pathway.**
M.G. Tansey, G.C. Chu, and J.P. Merlie

477 **Weaver granule neurons are rescued by calcium channel antagonists and antibodies against a neurite outgrowth domain of the B2 chain of laminin.**
P. Liesi and J.M. Wright

487 **Cutaneous overexpression of NT-3 increases sensory and sympathetic neuron number and enhances touch dome and hair follicle innervation.**
K.M. Albers, T.N. Perrone, T.P. Goodness, M.E. Jones, M.A. Green, and B.M. Davis

499 **Activity-induced internalization and rapid degradation of sodium channels in cultured fetal neurons.**
C. Paillart, J.-L. Boudier, J.-A. Boudier, H. Rochat, F. Couraud, and B. Dargent

511 **Biochemical properties and subcellular distribution of the BI and rbA isoforms of α_{1A} subunits of brain calcium channel.**
T. Sakurai, R.E. Westenbroek, J. Rettig, J. Hell, and W.A. Catterall

529 **Nuclear translocation of fibroblast growth factor (FGF) receptors in response to FGF-2.**
P.A. Maher

537 ***eat-5* and *unc-7* represent a multigene family in *Caenorhabditis elegans* involved in cell–cell coupling.**
T.A. Starich, R.Y.N. Lee, C. Panzarella, L. Avery, and J.E. Shaw

549 **Mechanism for transition from initial to stable cell–cell adhesion: Kinetic analysis of E-cadherin–mediated adhesion using a quantitative adhesion assay.**
B. Angres, A. Barth, and W.J. Nelson

559 **β4 integrin is required for hemidesmosome formation, cell adhesion and cell survival.**
J. Dowling, Q.-C. Yu, and E. Fuchs

573 **Altered rate of fibronectin matrix assembly by deletion of the first type III repeats.**
J.L. Sechler, Y. Takada, and J.E. Schwarzbauer

Volume 134, Number 3, August 1996

589 **Molecular and functional characterization of the p62 complex, an assembly of nuclear pore complex glycoproteins.**
T. Hu, T. Guan, and L. Gerace

603 **The cold sensitivity of a mutant of *Saccharomyces cerevisiae* lacking a mitochondrial heat shock protein 70 is suppressed by loss of mitochondrial DNA.**
B. Schilke, J. Forster, J. Davis, P. James, W. Walter, S. Laloraya, J. Johnson, B. Miao, and E. Craig

615 **Neither type of mannose 6-phosphate receptor is sufficient for targeting of lysosomal enzymes along intracellular routes.**
D. Kasper, F. Dittmer, K. von Figura, and R. Pohlmann

625 **The glucose transporter (GLUT-4) and vesicle-associated membrane protein-2 (VAMP-2) are segregated from recycling endosomes in insulin-sensitive cells.**
S. Martin, J. Tellam, C. Livingstone, J.W. Slot, G.W. Gould, and D.E. James

637 **ZP3-dependent activation of sperm cation channels regulates acrosomal secretion during mammalian fertilization.**
C. Arnoult, Y. Zeng, and H.M. Florman

647 **Lipid-modified, cysteinyl-containing peptides of diverse structures are efficiently S-acylated at the plasma membrane of mammalian cells.**
H. Schroeder, R. Leventis, S. Shahinian, P.A. Walton, and J.R. Silvius

661 **Ubiquitination of the yeast α-factor receptor.**
A.F. Roth and N.G. Davis

675 **Xenopus nonmuscle myosin heavy chain isoforms have different subcellular localizations and enzymatic activities.**
C.A. Kelley, J.R. Sellers, D.L. Gard, D. Bui, R.S. Adelstein, and I.C. Baines

689 **A role for gelsolin in actuating epidermal growth factor receptor–mediated cell motility.**
P. Chen, J.E. Murphy-Ullrich, and A. Wells

699 ***emo-1*, a *Caenorhabditis elegans* Sec61p γ homologue, is required for oocyte development and ovulation.**
K. Iwasaki, J. McCarter, R. Francis, and T. Schedl

715 **Envoplakin, a novel precursor of the cornified envelope that has homology to desmoplakin.**
C. Ruhrberg, M.A.N. Hajibagheri, M. Simon, T.P. Dooley, and F.M. Watt

731 **Pantophysin is a ubiquitously expressed synaptophysin homologue and defines constitutive transport vesicles.**
N.K. Haass, J. Kartenbeck, and R.E. Leube

747 **Regulation of tyrosinase gene expression by cAMP in B16 melanoma cells involves two CATGTG motifs surrounding the TATA box: Implication of the microphthalmia gene product.**
C. Bertolotto, K. Bille, J.-P. Ortonne, and R. Ballotti

757 **Induction of neuronal apoptosis by camptothecin, an inhibitor of DNA topoisomerase-I: Evidence for cell cycle–independent toxicity.**
E.J. Morris and H.M. Geller

771 **Identification of stromal cell products that interact with pre-B cells.**
K. Oritani and P.W. Kincade

783 **FGF-1–dependent proliferative and migratory responses are impaired in senescent human umbilical vein endothelial cells and correlate with the inability to signal tyrosine phosphorylation of fibroblast growth factor receptor-1 substrates.**
S. Garfinkel, X. Hu, I.A. Prudovsky, G.A. McMahon, E.M. Kapnik, S.D. McDowell, and T. Maciag

793 **Control of adhesion-dependent cell survival by focal adhesion kinase.**
S.M. Frisch, K. Vuori, E. Ruoslahti, and P.-Y. Chan-Hui

801 **Regulated binding of a PTP1B-like phosphatase to N-cadherin: Control of cadherin-mediated adhesion by dephosphorylation of β-catenin.**
J. Balsamo, T. Leung, H. Ernst, M.K.B. Zanin, S. Hoffman, and J. Lilien

Volume 134, Number 4, August 1996

815 **The periphery of nuclear domain 10 (ND10) as site of DNA virus deposition.**
A.M. Ishov and G.G. Maul

827 **Regulation of RARβ2 mRNA expression: Evidence for an inhibitory peptide encoded in the 5′-untranslated region.**
K. Reynolds, A.M. Zimmer, and A. Zimmer

837 **Gene trapping in differentiating cell lines: Regulation of the lysosomal protease cathepsin B in skeletal myoblast growth and fusion.**
J.A. Gogos, R. Thompson, W. Lowry, B.F. Sloane, H. Weintraub, and M. Horwitz

849 **Targeting of human catalase to peroxisomes is dependent upon a novel COOH-terminal peroxisomal targeting sequence.**
P.E. Purdue and P.B. Lazarow

863 **Mechanisms of mutations inhibiting fusion and infection by Semliki forest virus.**
M. Kielian, M.R. Klimjack, S. Ghosh, and W.A. Duffus

873 **Transgenic *mdx* mice expressing dystrophin with a deletion in the actin-binding domain display a "mild becker" phenotype.**
K. Corrado, J.A. Rafael, P.L. Mills, N.M. Cole, J.A. Faulkner, K. Wang, and J.S. Chamberlain

885 ***unc-68* encodes a ryanodine receptor involved in regulating *C. elegans* body-wall muscle contraction.**
E.B. Maryon, R. Coronado, and P. Anderson

895 **Glycine 699 is pivotal for the motor activity of skeletal muscle myosin.**
F. Kinose, S.X. Wang, U.S. Kidambi, C.L. Moncman and D.A. Winkelmann

911 **Chemoattractant-mediated increases in cGMP induce changes in *Dictyostelium* myosin II heavy chain–specific protein kinase C activities.**
A. Dembinsky, H. Rubin, and S. Ravid

923 **Isolation and characterization of a *Drosophila* gene essential for early embryonic development and formation of cortical cleavage furrows.**
C.X. Zhang, M.P. Lee, A.D. Chen, S.D. Brown, and T.-S. Hsieh

935 **Aluminum fluoride stimulates surface protrusions in cells overexpressing the ARF6 GTPase.**
H. Radhakrishna, R.D. Klausner, and J.G. Donaldson

949 **Aberrant mitosis in fission yeast mutants defective in fatty acid synthetase and acetyl CoA carboxylase.**
S. Saitoh, K. Takahashi, K. Nabeshima, Y. Yamashita, Y. Nakaseko, A. Hirata, and M. Yanagida

963 **Role of inhibitory CDC2 phosphorylation in radiation-induced G2 arrest in human cells.**
P. Jin, Y. Gu, and D.O. Morgan

971 **The function of intermediate filaments in cell shape and cytoskeletal integrity.**
R.D. Goldman, S. Khuon, Y.H. Chou, P. Opal, and P.M. Steinert

985 **Breaking the connection: Displacement of the desmosomal plaque protein desmoplakin from cell–cell interfaces disrupts anchorage of intermediate filament bundles and alters intercellular junction assembly.**
E.A. Bornslaeger, C.M. Corcoran, T.S. Stappenbeck, and K.J. Green

1003 **Symplekin, a novel type of tight junction plaque protein.**
B.H. Keon, S. Schäfer, C. Kuhn, C. Grund, and W.W. Franke

1019 **Properties and regulation of gap junctional hemichannels in the plasma membranes of cultured cells.**
H. Li, T.-F. Liu, A. Lazrak, C. Peracchia, G.S. Goldberg, P.D. Lampe, and R.G. Johnson

1031 **Functional dissociation of paracellular permeability and transepithelial electrical resistance and disruption of the apical–basolateral intramembrane diffusion barrier by expression of a mutant tight junction membrane protein.**
M.S. Balda, J.A. Whitney, C. Flores, S. González, M. Cereijido, and K. Matter

1051 **A novel calmodulin-binding protein, belonging to the WD-repeat family, is localized in dendrites of a subset of CNS neurons.**
F. Castets, M. Bartoli, J.V. Barnier, G. Baillat, P. Salin, A. Moqrich, J.-P. Bourgeois, F. Denizot, G. Rougon, G. Calothy, and A. Monneron

1063 **Sequential regulation of α4β1 and α5β1 integrin avidity by CC chemokines in monocytes: Implications for transendothelial chemotaxis.**
C. Weber, R. Alon, B. Moser, and T.A. Springer

1075 **CD44 and hyaluronan-dependent rolling interactions of lymphocytes on tonsillar stroma.**
R.A. Clark, R. Alon, and T.A. Springer

1089 **Early detachment of colon carcinoma cells during CD95(APO-1/Fas)-mediated apoptosis I. De-adhesion from hyaluronate by shedding of CD44.**
A.R. Günthert, J. Sträter, U. von Reyher, C. Henne, S. Joos, K. Koretz, G. Moldenhauer, P.H. Krammer, and P. Möller

---

Volume 134, Number 5, September 1996

1097 **Topoisomerase II α is associated with the mammalian centromere in a cell cycle– and species-specific manner and is required for proper centromere/kinetochore structure.**
J.B. Rattner, M.J. Hendzel, C.S. Furbee, M.T. Muller, and D.P. Bazett-Jones

1109 **Centromere and telomere movements during early meiotic prophase of mouse and man are associated with the onset of chromosome pairing.**
H. Scherthan, S. Weich, H. Schwegler, C. Heyting, M. Härle, and T. Cremer

1127 **Bipolar spindle attachments affect redistributions of ZW10, a *Drosophila* centromere/kinetochore component required for accurate chromosome segregation.**
B.C. Williams, M. Gatti, and M.L. Goldberg

1141 **Targeting and function in mRNA export of nuclear pore complex protein Nup153.**
R. Bastos, A. Lin, M. Enarson, and B. Burke

1157 **A nuclear export signal is essential for the cytosolic localization of the ran binding protein, RanBP1.**
S.A. Richards, K.M. Lounsbury, K.L. Carey, and I.G. Macara

1169 **Clustered folate receptors deliver 5-methyltetrahydrofolate to cytoplasm of MA104 cells.**
E.J. Smart, C. Mineo, and R.G.W. Anderson

1179 **Two muscle-specific LIM proteins in *Drosophila*.**
B.E. Stronach, S.E. Siegrist, and M.C. Beckerle

1197 **Talin and vinculin play distinct roles in filopodial motility in the neuronal growth cone.**
A.M. Sydor, A.L. Su, F.-S. Wang, A. Xu, and D.G. Jay

1209 **Coordination of protrusion and translocation of the keratocyte involves rolling of the cell body.**
K.I. Anderson, Y.-L. Wang, and J.V. Small

1219 **Synaptic vesicle recycling in synapsin I knock-out mice.**
T.A. Ryan, L. Li, L.-S. Chin, P. Greengard, and S.J Smith

1229 **Targeting of P-selectin to two regulated secretory organelles in PC12 cells.**
J.P. Norcott, R. Solari, and D.F. Cutler

1241 **Accumulation in fetal muscle and localization to the neuromuscular junction of cAMP-dependent protein kinase A regulatory and catalytic subunits RIα and Cα.**
T. Imaizumi-Scherrer, D.M. Faust, J.-C. Bénichou, R. Hellio, and M.C. Weiss

1255 **Disruption of muscle architecture and myocardial degeneration in mice lacking desmin.**
D.J. Milner, G. Weitzer, D. Tran, A. Bradley, and Y. Capetanaki.

1271 **β-Catenin associates with the actin-bundling protein fascin in a noncadherin complex.**
Y.S. Tao, R.A. Edwards, B. Tubb, S. Wang, J. Bryan, and P.D. McCrea

1283 **An in vivo structure-function study of armadillo, the β-catenin homologue, reveals both separate and overlapping regions of the protein required for cell adhesion and for wingless signaling.**
S. Orsulic and M. Peifer

1301 **PDGF induction of α_2 integrin gene expression is mediated by protein kinase C-ζ.**
J. Xu, M.M. Zutter, S.A. Santoro, and R.A.F. Clark

1313 **Integrin-associated protein immunoglobulin domain is necessary for efficient vitronectin bead binding.**
F.P. Lindberg, H.D. Gresham, M.I. Reinhold, and E.J. Brown

1323 **Protein kinase C regulates αvβ5-dependent cytoskeletal associations and focal adhesion kinase phosphorylation.**
J.M. Lewis, D.A. Cheresh, and M.A. Schwartz

1333 **A potent far-upstream enhancer in the mouse Proα2(I) collagen gene regulates expression of reporter genes in transgenic mice.**
G. Bou-Gharios, L.A. Garrett, J. Rossert, K. Niederreither, H. Eberspaecher, C. Smith, C. Black, and B. de Crombrugghe

Mini-review

1345 **Septins may form a ubiquitous family of cytoskeletal filaments.**
J.A. Cooper and D.P. Kiehart

Regular Articles

1349 **The clustering of telomeres and colocalization with Rap1, Sir3, and Sir4 proteins in wild-type *Saccharomyces cerevisiae.***
M. Gotta, T. Laroche, A. Formenton, L. Maillet, H. Scherthan, and S.M. Gasser

1365 **The hnRNP C proteins contain a nuclear retention sequence that can override nuclear export signals.**
S. Nakielny and G. Dreyfuss

1375 **The molecular chaperone Hsp78 confers compartment-specific thermotolerance to mitochondria.**
M. Schmitt, W. Neupert, and T. Langer

1387 **Dynamic measurement of the pH of the Golgi complex in living cells using retrograde transport of the verotoxin receptor.**
J.H. Kim, C.A. Lingwood, D.B. Williams, W. Furuya, M.F. Manolson, and S. Grinstein

1401 **Protein kinase C μ is located at the Golgi compartment.**
J. Prestle, K. Pfizenmaier, J. Brenner, and F.-J. Johannes

1411 **Sorting by COP I–coated vesicles under interphase and mitotic conditions.**
B. Sönnichsen, R. Watson, H. Clausen, T. Misteli, and G. Warren

1427 **FcεRI-mediated association of 6-μm beads with RBL-2H3 mast cells results in exclusion of signaling proteins from the forming phagosome and abrogation of normal downstream signaling.**
L. Pierini, D. Holowka, and B. Baird

1441 **The structure of the sarcomeric M band: Localization of defined domains of myomesin, M-protein, and the 250-kD carboxy-terminal region of titin by immunoelectron microscopy.**
W.M.J. Obermann, M. Gautel, F. Steiner, P.F.M. van der Ven, K. Weber, and D.O. Fürst

1455 **Basic amino acid residue cluster within nuclear targeting sequence motif is essential for cytoplasmic plectin-vimentin network junctions.**
B. Nikolic, E. Mac Nulty, B. Mir, and G. Wiche

1469 **Dlg protein is required for junction structure, cell polarity, and proliferation control in *Drosophila* epithelia.**
D.F. Woods, C. Hough, D. Peel, G. Callaini, and P.J. Bryant

1483 **Merosin and laminin in myogenesis; specific requirement for merosin in myotube stability and survival.**
P.H. Vachon, F. Loechel, H. Xu, U.M. Wewer, and E. Engvall

1499 **Tenascin-Y: A protein of novel domain structure is secreted by differentiated fibroblasts of muscle connective tissue.**
C. Hagios, M. Koch, J. Spring, M. Chiquet, and R. Chiquet-Ehrismann

1513 **Lack of association between receptor protein tyrosine phosphatase RPTPμ and cadherins.**
G.C.M. Zondag, W.H. Moolenaar, and M.F.B.G. Gebbink

1519 **Association between a transmembrane protein tyrosine phosphatase and the cadherin–catenin complex.**
R.M. Kypta, H. Su, and L.F. Reichardt

1531 **Dominant-negative effect on adhesion by myelin Po protein truncated in its cytoplasmic domain.**
M.-H. Wong and M.T. Filbin

1543 **The cell density factor CMF regulates the chemoattractant receptor cAR1 in *Dictyostelium.***
P.J.M. Van Haastert, J.D. Bishop, and R.H. Gomer

1551 **Modulation of cell migration by integrin-mediated cytoskeletal linkages and ligand-binding affinity.**
A. Huttenlocher, M.H. Ginsberg, and A.F. Horwitz

1563 **Is plasminogen activator inhibitor-1 the molecular switch that governs urokinase receptor-mediated cell adhesion and release?**
G. Deng, S.A. Curriden, S. Wang, S. Rosenberg, and D.J. Loskutoff

1573 **Separable *cis*-regulatory elements that contribute to tissue- and site-specific α2(XI) collagen gene expression in the embryonic mouse cartilage.**
N. Tsumaki, T. Kimura, Y. Matsui, K. Nakata, and T. Ochi

1583 **AUTHOR INDEX FOR VOLUME 134**

1591 **SUBJECT INDEX FOR VOLUME 134**